

# R Code and Examples for “Simple” Linear Regression

20191004 – Sleuth 3 Chapter 7

## Example

We have a data set with information about 152 flights by Endeavour Airlines that departed from JFK airport in New York to either Nashville (BNA), Cincinnati (CVG), or Minneapolis-Saint Paul (MSP) in January 2012.

```
head(flights, 4)
```

```
## # A tibble: 4 x 3
##   distance air_time dest
##   <dbl>    <dbl> <chr>
## 1     1029      189 MSP
## 2      765      150 BNA
## 3     1029      173 MSP
## 4      589      118 CVG
```

Fit the model and print a summary:

```
model_fit <- lm(air_time ~ distance, data = flights)
summary(model_fit)
```

```
##
## Call:
## lm(formula = air_time ~ distance, data = flights)
##
## Residuals:
##   Min       1Q   Median       3Q      Max
## -20.022  -7.054  -1.086   6.170  24.170
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept) 14.567729   3.955477   3.683 0.000321 ***
## distance     0.146999   0.004372  33.624 < 2e-16 ***
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 9.881 on 150 degrees of freedom
## Multiple R-squared:  0.8829, Adjusted R-squared:  0.8821
## F-statistic: 1131 on 1 and 150 DF, p-value: < 2.2e-16
```

1. What model did we fit? (This model describes the relationships in the population.)

2. What is the equation describing the model’s estimated mean air time as a function of distance?

3. What is the estimated intercept and its interpretation?
4. Conduct a hypothesis test of the claim that when a flight travels 0 miles, its air time is 0 minutes.
5. What is the estimated slope and its interpretation?
6. Conduct a hypothesis test of the claim that a flight's air time is unrelated to the distance travelled.

**7. Conduct a hypothesis test of the claim that these planes are flying at an average speed that's the same as the typical cruising speed of commercial passenger aircraft.**

According to Wikipedia, the typical cruising speed of commercial passenger aircraft is about 560 miles per hour ([https://en.wikipedia.org/wiki/Cruise\\_\(aeronautics\)](https://en.wikipedia.org/wiki/Cruise_(aeronautics))). After some unit changes, this works out to about 0.107 minutes per mile.

```
# calculate t statistic  
(0.147 - 0.107) / 0.0044
```

```
## [1] 9.090909
```

```
# calculate 2-sided p-value  
# pt(-9.09, df = 152 - 2) finds the probability of getting a t statistic <= -9.09  
pt(-9.09, df = 152 - 2) + pt(9.09, df = 152 - 2, lower.tail = FALSE)
```

```
## [1] 5.49638e-16
```

**8. Find and interpret a 95% confidence interval for the slope of the line**

```
# automatic calculations  
confint(model_fit)
```

```
##           2.5 %    97.5 %  
## (Intercept) 6.7520812 22.3833778  
## distance    0.1383611  0.1556377
```

```
# manual calculations from the formula: get the multiplier for an individual 95% CI  
qt(0.975, df = 152 - 2)
```

```
## [1] 1.975905
```

```
# calculate lower and upper endpoints of confidence interval  
0.147 - 1.976 * 0.00437
```


```
## [1] 0.1383649
```

```
0.147 + 1.976 * 0.00437
```

```
## [1] 0.1556351
```

## R Code to make scatterplot with estimated line overlaid

```
ggplot(data = flights, mapping = aes(x = distance, y = air_time)) +  
  geom_point() +  
  geom_smooth(method = "lm", se = FALSE) +  
  theme_bw()
```



```
ggplot(data = flights, mapping = aes(x = distance, y = air_time)) +  
  geom_point() +  
  geom_smooth(method = "lm") +  
  theme_bw()
```

