


Warm Up: Motivation for the Neyman-Pearson Lemma

- Data Model: $X_1, \dots, X_5 \stackrel{\text{i.i.d.}}{\sim} \text{Normal}(\theta, 5^2)$
- Let's consider a test of the hypotheses $H_0 : \theta = 25$ vs. $H_A : \theta = 20$
- If H_0 is correct, then $\bar{X} \sim \text{Normal}(25, 5^2/5)$. If H_A is correct, then $\bar{X} \sim \text{Normal}(20, 5^2/5)$


1. The following pictures can be used to illustrate 3 different tests based on the sampling distribution of \bar{X} , all with $P(\text{Type I Error} \mid H_0 \text{ true}) = 0.05$. For each test,

- Shade in the area corresponding to the probability of a Type I Error
- Shade in the area corresponding to the power of the test


Test 1: Reject H_0 if $\bar{x} \leq 21.322$ (This is the likelihood ratio test: for values of $\bar{x} \leq 21.322$, the p-value is ≤ 0.05 .)



Test 2: Reject H_0 if $\bar{x} \leq 20.617$ or $\bar{x} \geq 29.383$



Test 3: Reject H_0 if $20.617 \leq \bar{x} \leq 21.322$ or $28.678 \leq \bar{x} \leq 29.383$



2. Which of the tests above has the highest power?

3. For the likelihood ratio test, write down how you would calculate the probability of making a Type I Error and the power of the test as suitable integrals of either $f_{\bar{X}|\theta}(\bar{x}|20)$ or $f_{\bar{X}|\theta}(\bar{x}|25)$. (You will have 1 integral for the probability of a Type I Error and a second for the power of the test.)