


## Warm Up: Power Functions for Hypothesis Tests

- Data Model:  $X_1, \dots, X_5 \stackrel{\text{i.i.d.}}{\sim} \text{Normal}(\theta, 5^2)$
- We saw that the likelihood ratio test is *equivalent* to a test based on  $\bar{x}$ . The p-value is  $P(\bar{X} \leq \bar{x} | \theta = 25)$  (“extreme” values of  $\bar{x}$  are those that are at least as small as  $\bar{x}$ )
- The *power* of the test is  $P(\text{reject } H_0 | H_0 \text{ incorrect})$


1. Consider a test of the hypotheses  $H_0 : \theta = 25$  vs.  $H_A : \theta = 20$ . Below is a picture showing the pdf  $f_{\bar{X}|\theta}(\bar{x}|20)$  of a  $\text{Normal}(20, 5^2/5)$  distribution and the pdf  $f_{\bar{X}|\theta}(\bar{x}|25)$  of a  $\text{Normal}(25, 5^2/5)$  distribution, along with a vertical line at  $q_5^{\text{null}}$ , the 5th percentile of the  $\text{Normal}(25, 5^2/5)$  distribution.

- Shade in the area corresponding to  $1 - \beta$ , the power of the likelihood ratio test if  $H_A$  is correct.
- Show how you would calculate the power of the test as an integral of either  $f_{\bar{X}|\theta}(\bar{x}|20)$  or  $f_{\bar{X}|\theta}(\bar{x}|25)$




2. Suppose that instead we were testing the hypotheses  $H_0 : \theta = 25$  vs.  $H_A : \theta = 22.5$ . Below is a picture showing the pdf  $f_{\bar{X}|\theta}(\bar{x}|22.5)$  of a  $\text{Normal}(22.5, 5^2/5)$  distribution and the pdf  $f_{\bar{X}|\theta}(\bar{x}|25)$  of a  $\text{Normal}(25, 5^2/5)$  distribution, along with a vertical line at  $q_5^{\text{null}}$ , the 5th percentile of the  $\text{Normal}(25, 5^2/5)$  distribution.

- Shade in the area corresponding to  $1 - \beta$ , the power of the likelihood ratio test if  $H_A$  is correct.
- Show how you would calculate the power of the test as an integral of either  $f_{\bar{X}|\theta}(\bar{x}|22.5)$  or  $f_{\bar{X}|\theta}(\bar{x}|25)$



3. Suppose that instead we were testing the hypotheses  $H_0 : \theta = 25$  vs.  $H_A : \theta = 17.5$ . Below is a picture showing the pdf  $f_{\bar{X}|\theta}(\bar{x}|17.5)$  of a Normal(17.5,  $5^2/5$ ) distribution and the pdf  $f_{\bar{X}|\theta}(\bar{x}|25)$  of a Normal(25,  $5^2/5$ ) distribution, along with a vertical line at  $q_5^{null}$ , the 5th percentile of the Normal(25,  $5^2/5$ ) distribution.

- Shade in the area corresponding to  $1 - \beta$ , the power of the likelihood ratio test if  $H_A$  is correct.
- Show how you would calculate the power of the test as an integral of either  $f_{\bar{X}|\theta}(\bar{x}|17.5)$  or  $f_{\bar{X}|\theta}(\bar{x}|25)$



4. For which of the alternative hypotheses above ( $\theta = 17.5$ ,  $\theta = 20$ , or  $\theta = 22.5$ ) is the power of the test largest? For which is the power smallest?